



Discovery Ion-Exchange SPE Products

Base Silica: Irregular shaped, acid washed; 50µm particle size, 70Å pore diameter, 480m²/g specific surface area, 0.9cm³/g pore volume

Discovery ion-exchange SPE products are specifically developed, tested and quality controlled for pharmaceutical and clinical applications. The Discovery ion-exchange product line offers excellent selectivity towards charged molecular species enabling the user to extract, isolate, purify, and concentrate charged ionizable pharmaceuticals (basic or acidic) from both polar and non-polar sample matrices.

- Switches sample matrices and protects the analytical column/instrument from unwanted sample matrix components
- Concentrates target analyte(s) and removes background interferences for increased sensitivity and accuracy
- Improve extraction selectivity through Discovery's narrower pore size distribution
- Acid washed to reduced metal chelating activity
- Excellent capacity for the baseline clean-up of solution phase combinatorial chemistry reactions (removing target molecules from reaction by-products and excess reagents)


A comprehensive line of ion-exchange chemistries (four different SPE chemistries) and hardware configurations (including 96-well plates) to meet your diverse and most demanding sample prep needs.

<p>DSC-NH₂</p> $\begin{array}{c} \\ \text{--- Si ---} \\ \end{array} \text{---} (\text{CH}_2)_3\text{NH}_2$ <p style="text-align: right; font-size: small;">G001631</p>	<ul style="list-style-type: none"> – Polymerically bonded, aminopropyl phase that is very polar in nature (hydrogen bonding) allowing for both normal phase and ion exchange applications – A weak anion exchanger with a pKa of 9.8. At pH 7.8 or below, the functional groups are positively charged – Allows the rapid release of very strong anions such as sulfonic acids that may be retained irreversibly on SAX (a quarternary amine sorbent that is always positively charged) – Can be used in some reversed phase applications (due to ethyl spacer); however, it is predominately used as an ion-exchanger or normal phase sorbent due to its polar nature
<p>DSC-SAX</p> $\begin{array}{c} \\ \text{--- Si ---} \\ \end{array} \text{---} (\text{CH}_2)_3\text{N}^+(\text{CH}_3)_3$ <p style="text-align: right; font-size: small;">G001629</p>	<ul style="list-style-type: none"> – A polymerically bonded quarternary amine that remains charged at all pH levels – Commonly used when extracting weaker cations (e.g., carboxylic acids) that may not bind strongly enough to weaker anion-exchangers – Selectivity can be modified by changing the counter ion with the appropriate buffer during conditioning
<p>DSC-WCX</p> $\begin{array}{c} \\ \text{--- Si ---} \\ \end{array} \text{---} (\text{CH}_2)_3\text{N}(\text{CH}_2\text{COONa})\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{COONa})_2$ <p style="text-align: right; font-size: small;">G001632</p>	<ul style="list-style-type: none"> – A polymerically bonded carboxy propyl phase with a Na counter ion and a pKa of 4.8 – Its weak cation exchange properties carries a negative charge at pH 6.8 or above – A pH of 2.8 or below neutralizes this phase for easier elution of strong cationic analytes that are neutralized only at extreme basic conditions – Typically used when dealing with very strong cationic (high pKa) compounds that may be irreversibly retained on strong cation exchangers
<p>DSC-SCX</p> $\begin{array}{c} \\ \text{--- Si ---} \\ \end{array} \text{---} (\text{CH}_2)_2 \text{---} \text{C}_6\text{H}_4 \text{---} \text{SO}_3^- \text{H}^+$ <p style="text-align: right; font-size: small;">G001630</p>	<ul style="list-style-type: none"> – A polymerically bonded, benzene sulfonic acid functional group with a H⁺ counter ion that is a strong cation exchanger due to its very low pKa (<1.0) – Silica support allows for use with very organic solvents (no shrinking/swelling) – Excellent capacity (0.8meq/g) for cleaning up solution phase combinatorial chemistry reactions (removing target molecules from reaction by-products and excess reagents) – The presence of the benzene ring offers some mixed-mode capabilities (hydrophobic interactions) that should be considered when extracting cations from aqueous matrices

Here's the Proof:

Competitor Comparison of 3-methylpyrazole and 4-methylpyrazole from Urine Using Discovery DSC-SCX

SPE Tube: Discovery DSC-SCX, 500mg/3mL
HPLC Column: Discovery C18, 15cm x 4.6mm, 5µm particle, preceded by a 2cm guard column and 0.5µm frit filter
Mobile Phase: MeOH:5mM phosphate buffer, pH 6 (20:80)
Flow Rate: 1mL/min
Temp.: 30°C
Det.: UV, 220nm
Inj.: 25µL diluted urine extract



Efficiency of Recovery

Compound (µg/mL)	Discovery DSC-SCX (n=3)		Leading Competitor SCX (n=2)	
	%Recovery	%RSD	%Recovery	%RSD
1. 3-methylpyrazole (1.0)	89.4	±10.2%	67.1	±20%
2. Aprobarbital (1.0)	79.4	±6.8%	50.5	±30%

SPE Procedure

1. Condition and Equilibrate with 2mL MeOH & 2mL DI water
2. Load 1mL urine sample spiked with 1µg/mL of each analyte
3. Wash w/ 2mL DI Water
4. Elute with 2mL 5% MeOH in 250mM phosphate buffer, pH 7.4
5. Quantify against external standards via HPLC analyses

Ordering Information:

Discovery Ion-Exchange SPE Products

Product	Qty./Pk	DSC-NH ₂	DSC-SAX	DSC-WCX	DSC-SCX
Discovery SPE Tubes					
50mg/1mL	108 qty/pk	52635-U	52661-U	52737-U	52684-U
100mg/1mL	108 qty/pk	52636-U	52662-U	52739-U	52685-U
500mg/3mL	54 qty/pk	52637-U	52664-U	52741-U	52686-U
500mg/6mL	30 qty/pk	52638-U	52665-U	52742-U	52688-U
1g/6mL	30 qty/pk	52640-U	52666-U	52743-U	52689-U
2g/12mL	30 qty/pk	52641-U	52677-U	52744-U	52690-U
5g/20mL	20 qty/pk	52642-U	52688-U	52745-U	52691-U
10g/60mL	20 qty/pk	52644-U	52699-U	52746-U	52692-U
Bulk packing	100g	57212-U	57214-U	57228-U	57221-U
Discovery SPE-96 Well Plates					
100mg/well	1 ea	575615-U	575618-U	575633-U	575621-U
50mg/well	1 ea	575616-U	575619-U	575634-U	575622-U
25mg/well	1 ea	575617-U	575620-U	575635-U	575623-U
Discovery Büchner Funnels					
55mm ID x 30mm H, 12.5g	6 qty/pk	Custom	Custom	Custom	Custom
70mm ID x 40mm H, 25g	6 qty/pk	Custom	Custom	Custom	Custom
90mm ID x 48mm H, 50g	6 qty/pk	Custom	Custom	Custom	Custom
110mm ID x 66mm H, 100g	6 qty/pk	Custom	Custom	Custom	Custom

Trademark

Discovery is a trademark of Sigma-Aldrich Co.

sigma-aldrich.com/supelco

Order/Customer Service 800-247-6628, 800-325-3010 • Fax 800-325-5052 • E-mail supelco@sial.com
 Technical Service 800-359-3041, 814-359-3041 • Fax 800-359-3044, 814-359-5468 • E-mail techservice@sial.com
 SUPELCO • Supelco Park, 595 North Harrison Road, Bellefonte, PA 16823-0048 • 814-359-3441



We are committed to the success of our Customers, Employees and Shareholders through leadership in Life Science, High Technology and Service.

The SIGMA-ALDRICH Family

© 2002 Sigma-Aldrich Co. Printed in USA. Supelco brand products are sold through Sigma-Aldrich, Inc. Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.



T402054
ESW